arXiv:1304.4458v1 [hep-ph] 16 Apr 2013

Hidden charmonium decays of Z_c in intermediate meson loops model

Gang Li*
Department of Physics, Qufu Normal University, Qufu 273165, China

(Dated: April 17, 2013)

The BESIII collaboration reported an observation of a charged charmonium-like structure $Z_c(3900)$ in the invariant mass spectrum of $J/\psi\pi^\pm$ in $e^+e^- \to J/\psi\pi^+\pi^-$ at the center of mass 4.260 MeV, which could be an analogue of $Z_b(10610)$ claimed by the Belle Collaboration. In this work, we investigate the hidden charmonium transitions of Z_c via intermediate $D\,D^*$ meson loops. Our results show that the intermediate $D\,D^*$ meson loops are crucial for driving the transitions of Z_c. We also make a predication for the hidden charmonium decays of Z_c's partner (Z_c').

PACS numbers: 13.25.Gv,14.40.Pq,13.75.Lb

I. INTRODUCTION

Recently, a new charged state $Z_c^\pm$ is observed in the $J/\psi\pi^\pm$ invariant mass spectrum of $Y(4260) \to J/\psi\pi^+\pi^-$ decay by the BESIII Collaboration [1]. The reported mass and width are $M_{Z_c^\pm} = 3899.0 \pm 3.6 \pm 4.9$MeV and $\Gamma_{Z_c^\pm} = 46 \pm 11.3 \pm 12.6$MeV [1]. Belle Collaboration also observed a new charged charmonium-like structure in the $J/\psi\pi^\pm$ invariant mass spectrum with 5.2σ significance, with mass $M_{Z_c^\pm} = 3894.5 \pm 6.6 \pm 4.5$MeV and width $\Gamma_{Z_c^\pm} = 63 \pm 24 \pm 26$MeV [2]. Differing from the other hidden charmonium-like states, such as $X(3872)$, $Y(4260)$ etc, Z_c is electric charged states. Such a state, if it exists, need at least four quarks as minimal constituents, which makes them ideal candidates for exotic hadrons beyond the conventional $q\bar{q}$ mesons. On the one hand, this chain decay mode reminds us of the observations of $Z_b(10610)$ and $Z_b'(10650)$ in $\Upsilon(5S) \to Z_b(Z_b')\pi \to \Upsilon(nS)\pi\pi$ [3, 4]. On the other hand, the mass of $Z_c(3900)$ is in the vicinity of $D\bar{D}^* + c.c.$. This similar phenomenon (mechanism) shows that $Z_c(3900)$ may be an analogue of $Z_b(10610)$ claimed by the Belle Collaboration. We also expect the experimental observation of Z_c' and the neutral partners.

Recently, many investigations have been carried out to explain this exotic states [5–12]. The results of Ref. [5] show that it is necessary to include explicit $Z_c(3900)$ poles, i.e. a resonance structure as an isovector partner of $X(3872)$ in order for a more detailed description of the data. Applying heavy quark spin symmetry and heavy flavour symmetry the authors found a promising isovector 1^{+-} $D\bar{D}^*$ virtual state near threshold that might very well be identified with the newly discovered $Z_c(3900)$ [1, 2]

The intermediate meson loop (IML) transitions, or known as final state interactions (FSI), have been one of the important non-perturbative transition mechanisms in many processes [13–27]. In the energy region of charmonium masses, with more and more data from Belle, BaBar, CLEO and BES-III, it is widely recognized that intermediate hadron loops may be closely related to a lot of non-perturbative phenomena observed in experiment [16–30], e.g. apparent OZI-rule violations, sizeable non-$D\bar{D}$ decay branching ratios for $\psi(3770)$, and the HSR violations in charmonium decays. Recently, the IML transitions are also applied to bottomnium decays [31–35]. By applying the on-shell approximation, the bottom meson loops were suggested to play an important role in the $\Upsilon(5S)$ transitions to the lower Υ states with the emission of two pions [31] or one η [32]. This mechanism seems to explain many unusual properties that make the $\Upsilon(5S)$ different from $\Upsilon(4S)$. Similar approach was also applied to the study of Z_b and Z_b' by Liu *et al.* [33]. Within a nonrelativistic effective field theory, the decays of the $Z_b(10610)$ and the $Z_b(10650)$ to $\Upsilon(nS)\pi$ and $h_b(mP)\pi$ are investigated in Ref. [34]. The power counting analysis in Ref. [34] shows that the triangle transition $Z_b \to h_b(mP)\pi$ is not suppressed compared to $Z_b \to \Upsilon(nS)\pi$, although the decay is via a P-wave. In Ref. [35], we investigated the transitions from the $Z_b(10610)$ and $Z_b(10650)$ to bottomonium states with emission of a pion via intermediate BB^* meson loops in the ELA. By investigating the α-dependence of

---

*Electronic address: gli@mail.qfnu.edu.cn

partial decay widths and ratios between different decay channels, we show that the intermediate BB^* meson loops are crucial for driving the transitions of $Z_b/Z_b' \to \Upsilon(nS)\pi$ with $n = 1,2,3$, and $h_b(mP)\pi$ with $m = 1$ and 2.

Since the $Z_c(3900)$ is very close to the $D\bar{D}^*$ threshold, the IML should be an possible mechanism in its decays. In this work, we will investigate the decays of $Z_c^{(\prime)} \to J/\psi\pi$, $\psi'\pi$ and $h_c\pi$ via intermediate D-meson loops in an effective Lagrangian approach (ELA) with quantum numbers $I^G(J^{PC}) = 1^+(1^{+-})$ for the Z_c/Z_c'. The paper is organized as follows. In Sec. II, we will introduce the formulae for the ELA. In Sec. III, the numerical results are presented. The Summary will be given in Sec. IV.



FIG. 1: The hadron-level diagrams for $Z_c^+ \to \psi\pi^+$ with $D^{(*)}D^{(*)}$ as the intermediate states. Similar diagrams for Z_c^- and Z_c^0 states decays.



FIG. 2: The hadron-level diagrams for $Z_c^+ \to h_c\pi^+$ with $D^{(*)}D^{(*)}$ as the intermediate states. Similar diagrams for Z_c^- and Z_c^0 states decays.

II. TRANSITION AMPLITUDE

In order to calculate the leading contributions from the bottomed meson loops, we need the leading order effective Lagrangians for the couplings. Based on the heavy quark symmetry and chiral symmetry [36, 37], the relevant effective Lagrangians used in this work are as follows,

$$\begin{aligned}
\mathcal{L}_{\psi D^{(*)}D^{(*)}} &= ig_{\psi DD}\psi_\mu(\partial^\mu D\bar{D} - D\partial^\mu\bar{D}) - g_{\psi D^*D}\varepsilon^{\mu\nu\alpha\beta}\partial_\mu\psi_\nu(\partial_\alpha D^*_\beta\bar{D} + D\partial_\alpha\bar{D}^*_\beta) \\
&\quad -ig_{\psi D^*D^*}\left\{\psi^\mu(\partial_\mu D^{*\nu}\bar{D}^*_\nu - D^{*\nu}\partial_\mu\bar{D}^*_\nu) + (\partial_\mu\psi_\nu D^{*\nu} - \psi_\nu\partial_\mu D^{*\nu})\bar{D}^{*\mu} + D^{*\mu}(\psi^\nu\partial_\mu\bar{D}^*_\nu - \partial_\mu\psi^\nu\bar{D}^*_\nu)\right\}, \quad (1)\\
\mathcal{L}_{h_cD^{(*)}D^{(*)}} &= g_{h_cD^*D}h_c^\mu(D\bar{D}^*_\mu + D^*_\mu\bar{D}) + ig_{h_cD^*D^*}\varepsilon^{\mu\nu\alpha\beta}\partial_\mu h_{c\nu}D^*_\alpha\bar{D}^*_\beta\,, \quad (2)\\
\mathcal{L}_{D^*D^{(*)}\pi} &= -ig_{D^*D\pi}(D_i\partial_\mu P_{ij}\bar{D}^{*\mu}_j - D^{*\mu}_i\partial_\mu P_{ij}\bar{D}_j) + \frac{1}{2}g_{D^*D^*\pi}\varepsilon^{\mu\nu\alpha\beta}D^*_{i\mu}\partial_\nu P_{ij}\overleftrightarrow{\partial}_\alpha\bar{D}^*_{j\beta}, \quad (3)\\
\mathcal{L}_{Z_c^{(\prime)}D^{(*)}D^{(*)}} &= g_{Z_c^{(\prime)}DD^*D}Z_c^{(\prime)\mu}(D\bar{D}^*_\mu + D^*_\mu\bar{D}) + ig_{Z_c^{(\prime)}D^*D^*}\varepsilon^{\mu\nu\alpha\beta}\partial_\mu Z^{(\prime)}_{c\nu}D^*_\alpha\bar{D}^*_\beta\,, \quad (4)
\end{aligned}$$

where $D^{(*)} = \left(D^{(*)+}, D^{(*)0}\right)$ and $\bar{D}^{(*)T} = \left(D^{(*)-}, \bar{D}^{(*)0}\right)$ correspond to the charmed meson isodoublets. The coupling constants will be determined in the following.

The loop transition amplitudes for the transitions in Figs. 1 and 2 can be expressed in a general form in the effective Lagrangian approach as follows:

$$M_{fi} = \int \frac{d^4q_2}{(2\pi)^4}\sum_{D^*\ \text{pol.}}\frac{T_1T_2T_3}{a_1a_2a_3}\prod_i\mathcal{F}_i(m_i, q_i^2) \quad (5)$$

where T_i $(i = 1,2,3)$ are the vertex functions; $a_i = q_i^2 - m_i^2$ $(i = 1,2,3)$ are the denominators of the intermediate meson propagators. We adopt the form factor, $\prod_i \mathcal{F}_i(m_i, q_i^2)$, which is a product of monopole form factors for each internal mesons, i.e.

$$\prod_i \mathcal{F}_i(m_i, q_i^2) \equiv \mathcal{F}_1(m_1, q_1^2)\mathcal{F}_2(m_2, q_2^2)\mathcal{F}_3(m_3, q_3^2) \ , \tag{6}$$

with

$$\mathcal{F}_i(m_i, q_i^2) \equiv \left(\frac{\Lambda_i^2 - m_i^2}{\Lambda_i^2 - q_i^2}\right) , \tag{7}$$

where $\Lambda_i \equiv m_i + \alpha\Lambda_{\rm QCD}$ and the QCD energy scale $\Lambda_{\rm QCD} = 220$ MeV. This form factor is supposed to parameterize the non-local effects of the vertex functions and remove the loop integral divergence.

The explicit transition amplitudes for $Z_c(p_i) \to D^{(*)}(q_1)D^{(*)}(q_3)[D^{(*)}(q_2)] \to \psi(p_f)\pi(p_\pi)$ via those triangle loops are given as follows:

$$\begin{aligned}
M_{DD^*[D]} &= (i)^3 \int \frac{d^4 q_2}{(2\pi)^4} [g_{Z_b D^* D} \varepsilon_{i\mu}][g_{\psi DD} \varepsilon_f^{*\rho} (q_1 - q_2)_\rho][g_{D^* D\pi} p_{\pi\theta}] \\
&\quad \frac{i}{q_1^2 - m_1^2} \frac{i}{q_2^2 - m_2^2} \frac{i(-g^{\mu\theta} + q_3^\mu q_3^\theta / m_3^2)}{q_3^2 - m_3^2} \prod_i \mathcal{F}_i(m_i, q_i^2) && (8) \\
M_{DD^*[D^*]} &= (i)^3 \int \frac{d^4 q_2}{(2\pi)^4} [g_{Z_c D^* D} \varepsilon_{i\mu}][g_{\psi D^* D} \varepsilon_{\rho\sigma\xi\tau} p_f^\rho \varepsilon_f^{*\sigma} q_2^\xi][-g_{D^* D^* \pi} \varepsilon_{\theta\phi\kappa\lambda} p_\pi^\kappa q_2^\lambda] \\
&\quad \times \frac{i}{q_1^2 - m_1^2} \frac{i(-g^{\tau\theta} + q_2^\tau q_2^\theta / m_2^2)}{q_2^2 - m_2^2} \frac{i(-g^{\mu\phi} + q_3^\mu q_3^\phi / m_3^2)}{q_3^2 - m_3^2} \prod_i \mathcal{F}_i(m_i, q_i^2) && (9) \\
M_{D^* D[D^*]} &= (i)^3 \int \frac{d^4 q_2}{(2\pi)^4} [g_{Z_c D^* D} \varepsilon_{i\mu}][g_{\psi D^* D^*} (g_{\rho\sigma} g_{\xi\tau} - g_{\rho\tau} g_{\sigma\xi} + g_{\rho\xi} g_{\sigma\tau}) \varepsilon_f^{*\rho} (q_1 + q_2)^\tau][-g_{B^* B\pi} p_{\pi\theta}] \\
&\quad \times \frac{i(-g^{\mu\xi} + q_1^\mu q_1^\xi / m_1^2)}{q_1^2 - m_1^2} \frac{i(-g^{\sigma\theta} + q_2^\sigma q_2^\theta / m_2^2)}{q_2^2 - m_2^2} \frac{i}{q_3^2 - m_3^2} \prod_i \mathcal{F}_i(m_i, q_i^2) && (10) \\
M_{D^* D^*[D]} &= (i)^3 \int \frac{d^4 q_2}{(2\pi)^4} [g_{Z_c D^* D^*} \varepsilon_{\mu\nu\alpha\beta} q_i^\mu \varepsilon_i^\nu][g_{\psi D^* D} \varepsilon_{\rho\sigma\xi\tau} p_f^\rho \varepsilon_f^{*\sigma} q_1^\xi][g_{D^* D\pi} p_{\pi\theta}] \\
&\quad \times \frac{i(-g^{\alpha\tau} + q_1^\alpha q_1^\tau / m_1^2)}{q_1^2 - m_1^2} \frac{i}{q_2^2 - m_2^2} \frac{i(-g^{\beta\theta} + q_3^\beta q_3^\theta / m_3^2)}{q_3^2 - m_3^2} \prod_i \mathcal{F}_i(m_i, q_i^2) && (11) \\
M_{D^* D^*[D^*]} &= (i)^3 \int \frac{d^4 q_2}{(2\pi)^4} [g_{Z_c D^* D^*} \varepsilon_{\mu\nu\alpha\beta} p_i^\mu \varepsilon_i^\nu][g_{\psi D^* D^*} (g_{\rho\sigma} g_{\xi\tau} - g_{\rho\tau} g_{\sigma\xi} + g_{\rho\xi} g_{\sigma\tau}) \varepsilon_f^{*\rho} (q_1 + q_2)^\tau][-g_{D^* D^* \pi} \varepsilon_{\theta\phi\kappa\lambda} p_\pi^\kappa q_2^\lambda] \\
&\quad \times \frac{i(-g^{\alpha\xi} + q_1^\alpha q_1^\xi / m_1^2)}{q_1^2 - m_1^2} \frac{i(-g^{\sigma\theta} + q_2^\sigma q_2^\theta / m_2^2)}{q_2^2 - m_2^2} \frac{i(-g^{\beta\phi} + q_3^\beta q_3^\phi / m_3^2)}{q_3^2 - m_3^2} \prod_i \mathcal{F}_i(m_i, q_i^2) \ . && (12)
\end{aligned}$$

Also, the explicit transition amplitudes for $Z_c(p_i) \to D^{(*)}(q_1)D^{(*)}(q_3)[D^{(*)}(q_2)] \to h_c(p_f)\pi(p_\pi)$ via those triangle



FIG. 3: (a) The α-dependence of the partial widths of $Z_c^+ \to J/\psi\pi^+$ (solid line) and $\psi'\pi^+$ (dashed line). (b) The α-dependence of the partial widths of $Z_c'^+ \to J/\psi\pi^+$ (solid line), $\psi'\pi^+$ (dashed line). The mass of Z_c' is fixed to be 4040 MeV.

loops are given as follows:

$$\begin{aligned} M_{DD^*[D^*]} &= (i)^3 \int \frac{d^4 q_2}{(2\pi)^4} [g_{Z_cD^*D}\varepsilon_{i\mu}][g_{h_cD^*D}\varepsilon^*_{f\rho}][-g_{D^*D^*\pi}\varepsilon_{\theta\phi\kappa\lambda}p_\pi^\kappa q_2^\lambda] \\ &\quad \times \frac{i}{q_1^2-m_1^2}\frac{i(-g^{\rho\theta}+q_2^\rho q_2^\theta/m_2^2)}{q_2^2-m_2^2}\frac{i(-g^{\mu\phi}+q_3^\mu q_3^\phi/m_3^2)}{q_3^2-m_3^2}\prod_i \mathcal{F}_i(m_i,q_i^2) \end{aligned} \tag{13}$$

$$\begin{aligned} M_{D^*D[D^*]} &= (i)^3 \int \frac{d^4 q_2}{(2\pi)^4} [g_{Z_cD^*D}\varepsilon_{i\mu}][g_{h_cD^*D^*}\varepsilon_{\rho\sigma\xi\tau}p_f^\rho\varepsilon_f^{*\sigma}][-g_{D^*D\pi}p_{\pi\theta}] \\ &\quad \times \frac{i(-g^{\mu\xi}+q_1^\mu q_1^\xi/m_1^2)}{q_1^2-m_1^2}\frac{i(-g^{\tau\theta}+q_2^\tau q_2^\theta/m_2^2)}{q_2^2-m_2^2}\frac{i}{q_3^2-m_3^2}\prod_i \mathcal{F}_i(m_i,q_i^2) \end{aligned} \tag{14}$$

$$\begin{aligned} M_{D^*D^*[D]} &= (i)^3 \int \frac{d^4 q_2}{(2\pi)^4} [g_{Z_cD^*D^*}\varepsilon_{\mu\nu\alpha\beta}p_i^\mu\varepsilon_0^\nu][g_{h_cD^*D}\varepsilon^*_{f\rho}][g_{D^*D\pi}p_{\pi\theta}] \\ &\quad \times \frac{i(-g^{\alpha\rho}+q_1^\alpha q_1^\rho/m_1^2)}{q_1^2-m_1^2}\frac{i}{q_2^2-m_2^2}\frac{i(-g^{\beta\theta}+q_3^\beta q_3^\theta/m_3^2)}{q_3^2-m_3^2}\prod_i \mathcal{F}_i(m_i,q_i^2) \end{aligned} \tag{15}$$

$$\begin{aligned} M_{D^*D^*[D^*]} &= (i)^3 \int \frac{d^4 q_2}{(2\pi)^4} [g_{Z_cD^*D^*}\varepsilon_{\mu\nu\alpha\beta}p_i^\mu\varepsilon_i^\nu][g_{h_cD^*D^*}\varepsilon_{\rho\sigma\xi\tau}p_f^\rho\varepsilon_f^{*\sigma}][-g_{D^*D^*\pi}\varepsilon_{\theta\phi\kappa\lambda}p_\pi^\kappa q_2^\lambda] \\ &\quad \times \frac{i(-g^{\alpha\xi}+q_1^\alpha q_1^\xi/m_1^2)}{q_1^2-m_1^2}\frac{i(-g^{\tau\theta}+q_2^\tau q_2^\theta/m_2^2)}{q_2^2-m_2^2}\frac{i(-g^{\beta\phi}+q_3^\beta q_3^\phi/m_3^2)}{q_3^2-m_3^2}\prod_i \mathcal{F}_i(m_i,q_i^2), \end{aligned} \tag{16}$$

where p_i, p_f, p_π are the four-vector momenta of the initial Z_c, final state charmonium and pion, respectively, and q_1, q_2, and q_3 are the four-vector momenta of the intermediate charmed mesons as defined in Figs. 1 and 2.

III. RESULTS

Before proceeding to the numerical results, we first discuss the parameters, such as the coupling constants, in the formulation given in Section. II. In Eq. (1), the following couplings are adopted in the numerical calculations,

$$g_{\psi DD} = 2g_2\sqrt{m_\psi}m_D\ , \quad g_{\psi D^*D} = \frac{g_{\psi DD}}{\sqrt{m_D m_{D^*}}}\ , \quad g_{\psi D^*D^*} = g_{\psi D^*D}\sqrt{\frac{m_{D^*}}{m_D}}m_{D^*}\ , \tag{17}$$

In principle, couplings g_2 must be computed by nonperturbative methods. It shows that vector meson dominance (VMD) would provide an estimate of these quantities [36]. The coupling g_2 can be related to the J/ψ leptonic



FIG. 4: (a) The α-dependence of the partial widths of $Z_c^+ \to h_c\pi^+$. (b) The α-dependence of the partial widths of $Z_c'^+ \to h_c\pi^+$. The mass of Z_c' is fixed to be 4040 MeV.



FIG. 5: (a) The partial widths of $Z_c^+ \to J/\psi\pi^+$ (solid line) and $\psi'\pi^+$ (dashed line) in terms of the mass of Z_c'. (b) The partial widths of $Z_c'^+ \to h_c\pi^+$ in terms of the mass of Z_c'. The cutoff parameter $\alpha = 0.8$.

constant f_ψ which is defined by the matrix element $\langle 0|\bar{c}\gamma_\mu c|J/\psi(p,\varepsilon)\rangle = f_\psi m_\psi \varepsilon^\mu$, and $g_2 = \sqrt{m_\psi}/2m_D f_\psi$, where $f_\psi = 405 \pm 14$ MeV, and we have applied the relation $g_{\psi DD} = m_\psi / f_\psi$.



FIG. 6: The α-dependence of the ratios R_{Z_c} (solid line) and $\mathrm{R}_{Z_c'}$ (dashed line) defined in Eq. (22).

The ratio of the coupling constants $g_{\psi' DD}$ to $g_{\psi DD}$ is fixed as Ref. [25]

$$\frac{g_{\psi' DD}}{g_{\psi DD}} = 0.9. \tag{18}$$

In addition, the coupling constants in Eq. (2) are determined as

$$g_{h_c DD^*} \;=\; -2g_1\sqrt{m_{h_c} m_D m_{D^*}}, \;\; g_{h_c D^* D^*} = 2g_1\frac{m_{D^*}}{\sqrt{m_{h_c}}}, \tag{19}$$

with $g_1 = -\sqrt{m_{\chi_{c0}}/3}/f_{\chi_{c0}}$, where $m_{\chi_{c0}}$ and $f_{\chi_{c0}}$ are the mass and decay constant of $\chi_{c0}(1P)$, respectively [38], i.e. $f_{\chi_{c0}} = 175 \pm 55$ MeV [39].

In analogy to what is known about the Z_b states we assume that the width of the $Z_c(3900)$ is saturated by the DD^* intermediate state. We obtain the coupling constant $g_{Z_c D^* D} = 6.85$ GeV. The relations

$$g_{Z_c' D^* D} = g_{Z_c D^* D}, \quad g_{Z_c^{(\prime)} D^* D^*} = -g_{Z_c^{(\prime)} D^* D}\sqrt{\frac{m_D^*}{m_D m^2_{Z_c^{(\prime)}}}} \tag{20}$$

are applied to extract the couplings for $g_{Z_c^{(\prime)} D^* D^*}$.

The coupling constants relevant to the pion interactions in Eq. (4) are

$$g_{D^* D\pi} = \frac{2g}{f_\pi}\sqrt{m_D m_{D^*}}\ , \quad g_{D^* D^* \pi} = \frac{g_{D^* D\pi}}{\sqrt{m_D m_{D^*}}}\ , \tag{21}$$

where f_π = 132 MeV is the pion decay constant, the parameters g_V respects the relation $g_V = m_\rho/f_\pi$ [37]. We take $\lambda = 0.56\,\mathrm{GeV}^{-1}$ and $g = 0.59$ [41].

In Fig. 3(a), we plot the α-dependence of the decay partial widths of $Z_c \to J/\psi\pi$ (solid line) and $\psi'\pi$ (dashed line), respectively. A predominant feature is that the α dependence of the partial widths are quite stable, which indicates a reasonable cutoff of the ultraviolet contributions by the empirical form factors. As shown in this figure, at the same α, the intermediate D-meson loop effects turn out to be more important in $Z_c \to \psi'\pi$ than in $Z_c \to J/\psi\pi$. As a result, a smaller value of α is favored in $Z_c \to \psi'\pi$. This is understandable since the mass of ψ' is closer to the thresholds of DD^* or D^*D^* than J/ψ [42]. Thus, it gives rise to important threshold effects in $Z_c \to \psi'\pi^+$.

One also notices that the α-dependence of the partial widths for $Z_c^+ \to \psi'\pi^+$ are stabler than that for $J/\psi\pi$. This indicates that the enhanced partial widths are not from the off-shell part of the loop integrals. As we know that the dispersive contributions become rather model-dependent near threshold, the enhanced (but rather stable in terms of α) partial widths for $Z_c^+ \to \psi'\pi^+$ suggest that more stringent dynamic constraints are presumably needed to describe the near-threshold phenomena where the local quark-hadron duality has been apparently violated. What makes this process different from e.g. $\psi' \to h_c\pi^0$ in Ref. [26] is that there is no cancelations between the charged and neutral meson loops. As a consequence, the subleading terms in Refs. [26, 27] become actually leading contributions.

For the Z_c' decays, we carried out the following two analysis to understand this exotic states due to the lack of experimental information for Z_c'. First, we fix the Z_c' mass and then study the cut-off parameter α dependence of $Z_c' \to J/\psi\pi$, $\psi'\pi$. Second, we fix the α values (Here we choose $\alpha = 0.8$) and then study the Z_c' mass dependence of these decays.

The measured central mass of Z_c is 23.7MeV above the $D^{*+}\bar{D}^0$ threshold and 22.2MeV above the $D^{*+}\bar{D}^0$ threshold [42]. So we put the Z_c' resonance by the same amount above the $D^{*+}\bar{D}^{*0}$ threshold which gives its expected mass at approximately 4040MeV [42]. In Fig. 3(b), we plot the α-dependence of the decay partial widths of $Z_c' \to J/\psi\pi$ (solid line) and $\psi'\pi$ (dashed line), respectively. For the $Z_c' \to J/\psi\pi$ and $\psi'\pi$, the situation becomes more complicated. The thresholds of intermediate meson, i.e., $m_{D^{(*)}} + m_{D^{(*)}} + 2\alpha\Lambda_{QCD}$, are very close to the initial Z_c' mass at small α region, which leads to much more dramatic dependence behavior. The behaviors become more stabler at relatively larger α region. It indicates that for sufficiently heavy decaying charmonia and small values of α these singularities are located nearby or even inside the physical regime and additionally introduce unphysical contributions. So a relatively larger α should be considered in order to avoid introduce unphysical contributions.

The α-dependence of the partial widths of $Z_c \to h_c\pi$ are shown in Fig. 4(a). From this figure, we can see that the intermediate meson loop contributions are important in $Z_c \to h_c\pi$. Again, the behavior of $Z_c \to h_c\pi$ is quite stable, which indicates the dominant mechanism driven by the intermediate meson loops. We make an prediction for the partial width of $Z_c' \to h_c\pi$, the results are plotted in FIg. 4(b) with $m_{Z_c'} = 4040$ MeV. Again, $Z_c' \to h_c\pi$ shows a similar behavior to that of $Z_c' \to J/\psi\pi$ and $\psi'\pi$.

For the Z_c' decay, we fix the $\alpha = 0.8$ and vary the mass of Z_c'. The predicated results are present in Fig. 5. As shown in this diagram, due to the thresholds effects, there is a peak near 4050 MeV.

It would be interesting to further clarify the uncertainties arising from the introduction of form factors by studying the α dependence of the ratios between different partial decay widths. For the decays of $Z_c^+/Z_c'^+ \to J/\psi\pi^+$, we define the following ratios to the partial decay widths of $Z_c^+/Z_c'^+ \to \psi'\pi^+$:

$$\mathrm{R}_{Z_c} = \frac{\Gamma(Z_c^+ \to J/\psi\pi^+)}{\Gamma(Z_c^+ \to \psi'\pi^+)}, \quad \mathrm{R}_{Z_c'} = \frac{\Gamma(Z_c'^+ \to J/\psi\pi^+)}{\Gamma(Z_c'^+ \to \psi'\pi^+)}, \tag{22}$$

which are plotted in Fig. 6. The stabilities of the ratios in terms of α indicate a reasonably controlled cutoff for each channels by the form factor. Since the first coupling vertices are the same for those decay channels when taking the ratio, the stability of the ratios suggests that the transitions of $Z_c/Z_c' \to J/\psi\pi$ and $\psi'\pi$ are largely driven by the open threshold effects via the intermediate D meson loops. Here, we should mention that since the mass of ψ' is closer to $D^{(*)}D^{(*)}$ thresholds than J/ψ, so the ratio only reflects the open threshold effects via the intermediate D meson loops. The future experimental measurements of $Z_c/Z_c' \to J/\psi\pi$ and $\psi'\pi$ can help us investigate this issue deeply.

In order to understand this, the following analysis is carried out. First, one notices that we have adopted the couplings for the h_c and ψ to $D\bar{D}^*$ or $D^*\bar{D}^*$ in the heavy quark approximation. Since the physical masses for D and D^* are adopted in the loop integrals, the form factor will introduce unphysical pole contributions of which the interferences with the nearby physical poles would lead to model-dependent uncertainties. By assuming $M_{D^*} = M_D = 1869$ MeV and $M_{D^*} = M_D = 2010$ MeV, namely, by partially restoring the local quark-hadron duality, we calculate the partial widths of Z_c^+ and $Z_c'^+ \to J/\psi\pi$, $\psi'\pi$ and $h_c\pi$. We expect that the partial restoration of the local quark-hadron duality will significantly lower the partial widths since there will be only one physical pole in the loop and the unphysical one can be easily isolated away from the physical one. This is a rather direct demonstration of the sensitivity of the meson loop behaviors when close to open threshold and when the dispersive part becomes dominant. It shows a relatively stable behavior, especially Z_c' decays at the small α region.

IV. SUMMARY

In this work, we investigate hidden-charm decays of the newly discovered resonances Z_c^+ via intermediate D-meson loops. In this calculation, the quantum numbers of the neutral partners of these two resonances are fixed to be $I^G(J^{PC}) = 1^+(1^{+-})$, which has the same favored quantum number of Z_b'. For Z_c decays, our results show that the α dependence of the partial widths are quite stable, which indicate the dominant mechanism driven by the intermediate meson loops with a fairly well control of the UV contributions. For the predicated Z_c' decays, there shows dramatic behavior at small α region, however, the behavior becomes quite stable at relatively lager α region. Our results show that it will become sensitive to the meson loop contributions when the final state mass threshold are close to the intermediate meson thresholds. Namely, the effects from the unphysical pole introduced by the form factors would interfere with the nearby physical poles from the internal propagators and lead to model-dependent uncertainties. It is also a consequence of the violation of local quark-hadron duality. Such a phenomenon has been discussed in Ref. [27]. We expect the experiments to search for the Z_c' resonance and the decays of $Z_c \to DD^*$ and $Z_c' \to D^*D^*$, which will help us investigate the $Z_c^{(\prime)}$ decays deeply.

Acknowlegements

Authors thank X.-H. Liu, Q. Wang and Prof. Q. Zhao for useful discussions. This work is supported, in part, by the National Natural Science Foundation of China (Grant Nos. 11035006, 11175104, and 11275113), the DFG and

NSFC joint project CRC 110, the Chinese Academy of Sciences (KJCX3-SYW-N2) and the Ministry of Science and Technology of China (2009CB825200).

---

[1] M. Ablikim *et al.* [BESIII Collaboration], arXiv:1303.5949 [hep-ex].
[2] Z. Q. Liu *et al.* [Belle Collaboration], arXiv:1304.0121 [hep-ex].
[3] I. Adachi [Belle Collaboration], arXiv:1105.4583 [hep-ex].
[4] A. Bondar *et al.* [Belle Collaboration], Phys. Rev. Lett. **108**, 122001 (2012) [arXiv:1110.2251 [hep-ex]].
[5] Q. Wang, C. Hanhart, Q. Zhao and , arXiv:1303.6355 [hep-ph].
[6] F. -K. Guo, C. Hidalgo-Duque, J. Nieves, M. P. Valderrama and , arXiv:1303.6608 [hep-ph].
[7] D. -Y. Chen, X. Liu, T. Matsuki and , arXiv:1303.6842 [hep-ph].
[8] R. Faccini, L. Maiani, F. Piccinini, A. Pilloni, A. D. Polosa, V. Riquer and , arXiv:1303.6857 [hep-ph].
[9] M. B. Voloshin, arXiv:1304.0380 [hep-ph].
[10] N. Mahajan, arXiv:1304.1301 [hep-ph].
[11] C. -Y. Cui, Y. -L. Liu, W. -B. Chen and M. -Q. Huang, arXiv:1304.1850 [hep-ph].
[12] E. Wilbring, H. -W. Hammer and U. -G. Mei?ner, arXiv:1304.2882 [hep-ph].
[13] X. Q. Li, D. V. Bugg and B. S. Zou, Phys. Rev. D **55**, 1421 (1997).
[14] Q. Zhao and B. S. Zou, Phys. Rev. D **74**, 114025 (2006) [arXiv:hep-ph/0606196].
[15] Q. Zhao, Phys. Lett. B **636**, 197 (2006) [arXiv:hep-ph/0602216].
[16] J. J. Wu, Q. Zhao and B. S. Zou, Phys. Rev. D **75**, 114012 (2007) [arXiv:0704.3652 [hep-ph]].
[17] X. Liu, X. Q. Zeng and X. Q. Li, Phys. Rev. D **74**, 074003 (2006) [arXiv:hep-ph/0606191].
[18] H. Y. Cheng, C. K. Chua and A. Soni, Phys. Rev. D **71**, 014030 (2005) [arXiv:hep-ph/0409317].
[19] V. V. Anisovich, D. V. Bugg, A. V. Sarantsev and B. S. Zou, Phys. Rev. D **51**, 4619 (1995).
[20] Q. Zhao, B. s. Zou and Z. b. Ma, Phys. Lett. B **631**, 22 (2005) [arXiv:hep-ph/0508088].
[21] G. Li, Q. Zhao and B. S. Zou, Phys. Rev. D **77**, 014010 (2008) [arXiv:0706.0384 [hep-ph]].
[22] Y. J. Zhang, G. Li and Q. Zhao, Phys. Rev. Lett. **102**, 172001 (2009) [arXiv:0902.1300 [hep-ph]].
[23] X. H. Liu and Q. Zhao, Phys. Rev. D **81**, 014017 (2010) [arXiv:0912.1508 [hep-ph]].
[24] X. H. Liu and Q. Zhao, J. Phys. G **38**, 035007 (2011) [arXiv:1004.0496 [hep-ph]].
[25] Q. Wang, X. -H. Liu and Q. Zhao, Phys. Lett. B **711**, 364 (2012) [arXiv:1202.3026 [hep-ph]].
[26] F. K. Guo, C. Hanhart, G. Li, U. G. Meissner and Q. Zhao, Phys. Rev. D **82**, 034025 (2010) [arXiv:1002.2712 [hep-ph]].
[27] F. K. Guo, C. Hanhart, G. Li, U. G. Meissner and Q. Zhao, Phys. Rev. D **83**, 034013 (2011) [arXiv:1008.3632 [hep-ph]].
[28] N. Brambilla, S. Eidelman, B. K. Heltsley, R. Vogt, G. T. Bodwin, E. Eichten, A. D. Frawley and A. B. Meyer *et al.*, Eur. Phys. J. C **71**, 1534 (2011) [arXiv:1010.5827 [hep-ph]].
[29] N. Brambilla *et al.* [Quarkonium Working Group Collaboration], hep-ph/0412158.
[30] N. Brambilla, A. Pineda, J. Soto and A. Vairo, Rev. Mod. Phys. **77**, 1423 (2005) [hep-ph/0410047].
[31] C. Meng and K. T. Chao, Phys. Rev. D **77**, 074003 (2008) [arXiv:0712.3595 [hep-ph]].
[32] C. Meng and K. T. Chao, Phys. Rev. D **78**, 074001 (2008) [arXiv:0806.3259 [hep-ph]].
[33] Z. -F. Sun, J. He, X. Liu, Z. -G. Luo and S. -L. Zhu, Phys. Rev. D **84**, 054002 (2011) [arXiv:1106.2968 [hep-ph]].
[34] M. Cleven, Q. Wang, F. -K. Guo, C. Hanhart, U. -G. Meissner, Q. Zhao and , arXiv:1301.6461 [hep-ph].
[35] G. Li, F. L. Shao, C. W. Zhao, Q. Zhao and , Phys. Rev. D **87**, 034020 (2013) arXiv:1212.3784 [hep-ph].
[36] P. Colangelo, F. De Fazio and T. N. Pham, Phys. Rev. D **69**, 054023 (2004) [arXiv:hep-ph/0310084].
[37] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio and G. Nardulli, Phys. Rept. **281**, 145 (1997) [arXiv:hep-ph/9605342].
[38] P. Colangelo, F. De Fazio and T. N. Pham, Phys. Lett. B **542**, 71 (2002) [hep-ph/0207061].
[39] E. V. Veliev, H. Sundu, K. Azizi and M. Bayar, Phys. Rev. D **82**, 056012 (2010) [arXiv:1003.0119 [hep-ph]].
[40] X. Li and M. B. Voloshin, arXiv:1207.2425 [hep-ph].
[41] C. Isola, M. Ladisa, G. Nardulli and P. Santorelli, Phys. Rev. D **68**, 114001 (2003) [arXiv:hep-ph/0307367].
[42] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).